

1
2
3 **Anxious attachment style predicts an enhanced cortisol response to group**
4
5 **psychosocial stress**
6
7

8
9 Smyth N*, Thorn L*, Oskis A[∞], Hucklebridge F*, Evans P*, Clow A*^α.
10
11

12
13 * Psychophysiology and Stress Research Group, Department of Psychology, University
14 of Westminster, 115 New Cavendish Street, London W1W 6UW, United Kingdom
15

16
17 [∞] Department of Psychology, Middlesex University, The Burroughs, London, NW4 4BT
18
19 United Kingdom
20
21

22
23 ^α Corresponding author: clowa@wmin.ac.uk; 0207 911 5000
24
25
26

27
28 **Running Head:** Stress reactivity and attachment style
29
30

31
32 **Key words:** Trier Social Stress Test; TSST-G; HPA axis; stress reactivity; saliva; group
33
34 stressor; healthy females
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Insecure attachment style is associated with poor health outcomes. A proposed pathway implicates the hypothalamic-pituitary-adrenal axis (HPA-axis), dysregulation of which is associated with a wide range of mental and physical ill-health. However data on stress reactivity in relation to attachment style is contradictory. This relationship was examined using the novel Trier Social Stress Test for groups (TSST-G): a group-based acute psychosocial stressor. Each participant, in the presence of other group members, individually performed public speaking and mental arithmetic tasks. Seventy-eight healthy young females (20.2 ± 3.2 years), in groups of up to 6 participants completed demographic information and the Vulnerable Attachment Style Questionnaire (VASQ), and were then exposed to the TSST-G. Physiological stress reactivity was assessed using salivary cortisol concentrations, measured on 7 occasions at 10-minute intervals. Vulnerable attachment predicted greater cortisol reactivity independent of age, smoking status, menstrual phase and BMI. Supplementary analysis indicated that insecure anxious attachment style (high scores on the insecurity and proximity seeking sub-scales of the VASQ) showed greater cortisol reactivity than participants with secure attachment style. Avoidant attachment style (high scores for insecurity and low scores for proximity seeking) was not significantly different from the secure attachment style. Attachment style was not associated with the timing of the cortisol peak or post-stress recovery in cortisol concentrations. These findings in healthy young females indicate subtle underlying changes in HPA axis function in relation to attachment style and may be important for future mental health and well-being.

Introduction

Attachment style is suggested to be important for regulating threat appraisal, stress response and recovery from stress, although the mechanisms underlying this complex interplay are not well understood (Diamond, 2001). Within the adult attachment literature, 2

1
2
3 insecure attachment style is generally conceptualised along two dimensions, namely
4 attachment anxiety and avoidance (Brennan et al., 1998). A securely attached individual is
5 considered to be an individual with low levels of both (Brennan et al., 1998). High attachment
6 anxiety is associated with preoccupation with the availability and responsiveness of the
7 other, maximization of negative experiences and hyper-vigilance to potential threat.
8

9
10
11 Attachment avoidance is associated with a tendency to devalue intimacy and dependency
12 and maximize autonomous behaviour strategies when faced with potential threat. Insecure
13 attachment style is known to predict a range of poor physical and mental health outcomes
14 (Bifulco et al., 2002a; Bifulco et al., 2002b; Carr et al., 2013; Jinyao et al., 2012; Puig et al.,
15 2013). The biological underpinnings of these links however are not clear.
16
17
18
19
20
21
22
23
24

25 One of the proposed pathways implicates the hypothalamic-pituitary-adrenal (HPA) axis
26 (Repetti et al., 2002), dysregulation of which is associated with a wide range of mental and
27 physical ill-health (McEwen, 2000). A flattened diurnal rhythm of cortisol secretion has been
28 reported in anxious attachment style (Oskis et al., 2011; Quirin et al., 2008) however there are
29 mixed findings from studies examining reactivity of the HPA axis in relation to attachment style.
30 For example in one study avoidant (but not anxious) attachment predicted enhanced stress-
31 induced cortisol responding in females (Powers et al., 2006) whilst the opposite was
32 found in another study (Quirin et al., 2008). Other studies have reported insecure dismissing
33 attachment style to predict enhanced cortisol reactivity (Pierrehumbert et al., 2012; Rifkin-
34 Graboi, 2008) whereas secure and dismissive attachment styles have been reported as
35 similar elsewhere (Kidd et al., 2011). Further studies show no relationship between cortisol
36 responding to a stressor and attachment style (e.g. Ditzen et al., 2008; Smeets, 2010).
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52 Whether attachment style predicts acute stress responding remains unclear. The disparity in
53 the literature may in part be related to the wide array of methodologies that have been used
54 to investigate this issue. Stressors have ranged from the Trier Social Stress test (TSST) for
55
56
57
58
59
60

1
2
3 individuals (e.g. Ditzen et al., 2008; Smeets, 2010) to experimental conflict negotiation (e.g.
4 Powers et al., 2006) visualization of hypothetical distressing situations (e.g. Rifkin-Graboi,
5 2008) and behavioural interference tasks (Kidd et al., 2011). The most commonly used tool
6
7 to assess attachment style in adult stress reactivity studies is the Experiences in Close
8 Relationships Scale (Brennan et al., 1995), which assesses attachment in romantic
9 relationships. The Vulnerable Attachment Style Questionnaire (VASQ; Bifulco et al., 2003) is
10 arguably a more appropriate measure for use in research investigating attachment and HPA
11 axis activity as, rather than romantic attachment, it focuses on how individuals generally
12 relate to others. Furthermore it performs somewhat better in predicting depression than other
13 self-report measures of attachment (Bifulco et al., 2003), as well as predicting negative
14 psychosocial well-being and mental health in university students (Carr et al., 2013). The
15 VASQ, was developed and validated in relation to an in-depth interview procedure
16 (Attachment Style Interview; Bifulco et al., 2002a; Bifulco et al., 2002b) which has been
17 used in previous research examining HPA axis activity and attachment style (Oskis et al.,
18 2014; Oskis et al., 2011).

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36 The Trier Social Stress Test (TSST; Kirschbaum et al., 1993) comprises uncontrollability and
37 socio-evaluative threat known to reliably activate HPA axis function (Dickerson et al., 2004)
38 and has recently been adapted for use in group settings: the TSST-G (von Dawans et al.,
39 2011). A primary motivation was to increase the rate of participant exposure to the TSST
40 but it provides the opportunity to examine the impact of social dynamics on stress reactivity
41 (Häusser et al., 2012). In the present study we adapted the TSST-G to maximise
42 opportunities for group interaction which may attenuate or increase stress reactivity
43 depending on the characteristics of the individual within the group. Given that those with high
44 attachment insecurity easily perceive threats in their environment, frequently experience
45 social interactions as stressful and excessively ruminate about psychologically distressing
46 experiences (Burnette et al., 2009; Shaver et al., 2002) they might find the group version of
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the TSST particularly stressful. We chose to investigate an all-female sample since sex is
4 known to moderate the link between attachment style and HPA reactivity (Kiecolt-Glaser et
5 al., 1996; Kirschbaum et al., 1995; Stroud et al., 2002). Young, healthy participants were
6 recruited to explore whether attachment style might be a pre-clinical indicator of vulnerability
7 rather than a consequence of concurrent poor health. Due to discrepancies in the stress
8 reactivity and attachment literature the aim of this study was to examine self-reported
9 attachment and physiological stress responding to a group psychosocial stressor.
10
11
12
13
14
15
16

17 **Methods**

20 *Participants*

21
22
23
24
25 Eighty-one female undergraduate student participants were recruited. They did not receive
26 financial incentives but did receive course credits. Cortisol data was missing for one
27 participant as the salivary volume was insufficient for assay purposes, and another
28 participant did not complete the attachment questionnaires. A single participant was
29 removed from the data set on the basis that their cortisol data were more than 5 standard
30 deviations above the mean for each sample, and their data remained as outliers following
31 square root transformation. Analyses were performed on 78 participants, age ranging from
32 18 to 33 (mean \pm SD: 20.1 \pm 3.1) years. Participants were ethnically diverse; of those who
33 disclosed their ethnicity, 26 were Asian (Indian, Chinese, Pakistani, Bangladeshi, Arabic), 31
34 were white European, 13 were African Caribbean, and 4 were mixed race.
35
36
37
38
39
40
41
42
43
44
45

46 To control for sex differences in cortisol reactivity only females were recruited. To reduce
47 the impact of variables known to influence cortisol reactivity exclusion criteria included
48 medication, illness and history of psychiatric illness. As cortisol reactivity is influenced by the
49 menstrual cycle and body mass index (BMI) the number of days since last period was
50 recorded, as was height and weight (Dockray et al., 2009; Smyth et al., 2013). Two
51
52
53
54
55
56

57 participants used oral contraceptives. The majority of participants (86%) were non-smokers. 5
58
59
60

Vulnerable Attachment Style Questionnaire (VASQ)

The VASQ (Bifulco et al., 2003) is a brief self-report tool, which assesses general adult attachment. It is designed to assess overall attachment vulnerability as well as two dimensions of attachment: a global dimension of attachment insecurity common to all insecure subtypes (representing a deep-rooted mistrust of others and their motives) and a proximity seeking dimension reflecting the strategy individuals use to manage their insecurity (i.e. some individuals with high insecurity develop excessive neediness and vigilance of others, whilst other individuals develop an aversion to closeness with others). The scale comprises 22 items measured on a five-point scale, ranging from strongly disagree to strongly agree. Low scores on the 12-item insecurity subscale (e.g. "I find it hard to trust others" and "People let me down a lot") represent secure attachment and high scores reflect insecure attachment. The proximity seeking subscale consists of 10 items (e.g. "I get anxious when people close to me are away" and "I look forward to spending time on my own"). Low scores represent propensity for avoidant behaviour and high scores reflect a need for closeness with others. Cronbach's alpha was .81 for the insecurity scale and .74 for the proximity scale. A total attachment vulnerability measure can be derived by summing items on both scales. The VASQ can also be used to categorise participants according to secure, insecure anxious or insecure avoidant attachment styles. The insecure anxious attachment style category is derived from high scores on both insecure and proximity-seeking measures. The insecure avoidant type category is derived from high scores on the insecurity scale and low scores on the proximity scale.

Procedure

The study was approved by the University of Westminster Ethics Committee. Following recruitment, groups of participants were invited to attend a test session at a set time and

1
2
3 place. In line with best practice guidelines (Kirschbaum et al., 1993; Smyth et al., 2013)
4
5 testing commenced in the afternoon between 13:00 and 15:00 hr, to control for changes in
6
7 basal cortisol secretion in the morning and following the post-prandial period. Participants
8
9 were asked to refrain from food, caffeine, alcohol, exercise and smoking 30 minutes prior to
10
11 the research session. The TSST-G (von Dawans et al., 2011) included 3 main phases: the
12
13 group preparatory period (30 min); the group stress task period (22 min); and a group resting
14
15 and debriefing period (40 min). During the preparatory period, groups of up to 6 participants
16
17 met in Room 1 where they were informally seated around a single table and introduced to
18
19 the experimenter, they were free to talk to each other at this time.
20
21

22
23 Following informed written consent, participants completed in silence demographic
24
25 questions, the date of their last menstruation and the VASQ, if they had not already completed
26
27 it on-line (it had been available since the beginning of the recruitment period). Each participant
28
29 then received a large sticker with a number between 1 and 6. They were informed that they
30
31 would be identified with this number during the task period and that the numbers would be
32
33 called in a random order. Participants were then introduced to the saliva-sampling method.
34
35 Following this participants were given 10-minute quiet time to prepare notes for a mock job
36
37 interview. They were asked to prepare a free 2-minute speech as if
38
39 applying for a job of their choice and to introduce themselves to the committee. They were
40
41 asked to convince the committee that they were the most suitable candidates for the
42
43 position. After this preparatory period the baseline saliva sample was collected immediately
44
45 prior to leaving Room 1. Participants were taken into Room 2 (a short distance away) and
46
47 instructed to stand in a straight line in front of the already seated committee, comprising one
48
49 woman and one man. The committee were wearing white laboratory coats and there were
50
51 two conspicuous video cameras pointing at the participants. A committee member called the
52
53 number of each participant in turn in a random order to make a 2-minute speech as if
54
55 applying for a job. After all participants gave their speech (a total of up to 12 minutes), the 7
56
57
58
59
60

1
2
3 committee asked the participants, in the same order, to serially subtract the number 17 from
4 a given number (e.g. 4878) as fast and accurately as possible for 80 seconds. Each
5 participant received an individual starting number to avoid learning effects. Standard
6 responses from the committee were followed where participants ended their speech before
7 the 2-minute duration (e.g. 'you still have time, please continue') or failed in the subtraction
8 task (e.g. 'you made a mistake please start again from the number Q') (von Dawans et al.,
9 2011). Immediately after all participants had completed the TSST-G, participants were
10 returned to Room 1, where they collected saliva samples every 10 minutes up to 40 minutes
11 following the TSST-G period. During this time they were debriefed.
12
13
14
15
16
17
18
19
20
21
22

23 *Saliva Sampling Collection*

24
25
26
27 Cortisol was measured in saliva samples collected using Salivettes (saliva sampling devices,
28 Sarstedt Ltd., Leicester, England) at baseline (immediately before the TSST-G: S1, at 0 min)
29 immediately after the public speaking task (S2, at 12 min), after the mental arithmetic task (S3,
30 at 22 min), and every 10 min up to 60 min (S4, at 32 min S5, at 42 min, S6, at 52 min, and S7,
31 at 62 min). This cortisol profile allowed us to capture the rise in cortisol, the cortisol peak, and
32 the decline of cortisol (i.e. the recovery period) (Dickerson et al., 2004; Smyth et
33 al., 2013). Saliva samples were frozen at -20°C until assayed at the University of
34 Westminster. Samples were thawed and centrifuged for 10 minutes at 3,500 rpm. Cortisol
35 concentrations were determined by enzyme linked immuno-sorbent assay developed by
36 Salimetrics LLC (USA). The standard range in the assay was 0.33–82.77 nmol/l. Intra and
37 inter-assay variations were both below 10%.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

51 *Statistical analysis*

1
2
3 Cortisol data were moderately skewed and therefore a square root transformation was
4 applied which normalised distributions, although cortisol concentrations shown in figures are
5 representative of original units. Descriptive statistics were explored for each cortisol sample
6 measured throughout the TSST-G procedure, and a one-way within-subjects analysis of
7 variance was conducted to examine differences in cortisol over time. Within subjects
8 contrasts were used to assess the pattern of cortisol secretion.
9
10
11
12
13
14
15
16

17 As participants performed the TSST-G tasks at slightly different time-points, cortisol
18 reactivity was computed for each individual as their peak sample minus baseline. Cortisol
19 recovery was computed as individual peak sample minus sample 7 (recovery). Pearson's
20 correlation coefficients were used to examine relationships between these cortisol indices,
21 VASQ attachment measures and demographics variables. Significant relationships between
22 cortisol and attachment measures were examined in a multiple regression analysis controlling
23 for variables known to affect cortisol stress reactivity. Participants were categorised according
24 to the VASQ attachment style: secure (n=20), insecure anxious (n=37) and insecure avoidant
25 (n=21). A one-way between-subjects analysis of variance explored group differences in cortisol
26 stress reactivity and Bonferroni post hoc tests were applied. Chi-square was used to examine
27 the association between participants' peak cortisol
28 time and attachment style group.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 **Results**

47
48
49 Results indicated that the TSST-G induced an overall cortisol response in this sample ($F_{(6, 462)} = 7.623, p < .001$), illustrated in Figure 1. Within subjects contrasts revealed a significant
50 quadratic effect ($F_{(1, 77)} = 23.807, p < .001$), such that on average cortisol increased from
51
52
53
54
55
56
57
58
59
60

1
2
3 baseline peaked at the fourth sampling point (10 min after the completion of the TSST-G)
4
5 and subsequently declined.
6
7

8
9 **Insert Figure 1 here**
10

11
12
13 Relationships between cortisol data and attachment variables measured by the VASQ were
14 examined using focused composite cortisol indices: individual peak sample minus baseline
15 (cortisol reactivity) and individual peak sample minus sample 7 (recovery). Descriptive
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

23 **Insert table 1 here**
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

27 There was a significant positive relationship between cortisol reactivity and VASQ
28 vulnerability score ($r = .289, p = .010$) in that participants with a higher level of vulnerable
29 attachment exhibited a greater increase in cortisol from baseline to peak value. In other
30 terms, participants who demonstrated an insecure anxious attachment style (those scoring
31 highly on both VASQ subscales) displayed greater cortisol reactivity. With regards to the
32 dimensions of the VASQ, insecurity was significantly positively correlated with cortisol
33 reactivity, whereas proximity was not. There were no relationships between attachment
34 measures and cortisol recovery. There were also no relationships between cortisol
35 measures and pertinent demographic characteristics, apart from age, which was positively
36 related with both cortisol reactivity and recovery. In terms of VASQ attachment measures,
37 vulnerability was unrelated to demographic variables, however, insecurity was positively
38 related with age, and proximity was positively correlated with smoking status.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

54 A multiple regression analysis was conducted to examine whether the relationship between
55 cortisol reactivity and vulnerable attachment remained significant when variables known to
56
57
58
59
60

1
2
3 affect cortisol reactivity were included in the model (Table 2). Vulnerable attachment and age
4 remained significant independent predictors of cortisol reactivity.
5
6
7

8
9 **Insert Table 2 here**
10

11
12
13 The VASQ can be used to group participants according to secure, anxious or avoidant
14 attachment styles. The insecure anxious attachment style category is derived from high
15 scores on both insecure and proximity-seeking measures. In a supplementary analysis a
16 one-way between subjects ANOVA was performed to examine the difference in cortisol
17 reactivity between the three groups. There was a significant effect of attachment style group
18 on cortisol reactivity ($F_{(2,75)} = 5.300, p = .007$), see Figure 2. Bonferroni post hoc tests
19 indicated that the insecure anxious group was significantly different from the secure group (p
20 = .011). There was no association between when participants peaked and their attachment
21 style group ($\chi^2 = 16.405, p = .173$).
22
23
24
25
26
27
28
29
30
31

32
33 **Insert Figure 2 here**
34
35

36 37 **Discussion**

38
39
40
41 Vulnerable attachment, determined by the VASQ, predicted greater cortisol reactivity to a
42 group psychosocial stressor independent of age, smoking status, menstrual phase and BMI.
43
44 Whilst there were no differences in the timing of the cortisol peak, supplementary analysis
45 revealed that participants with an insecure anxious attachment style (a combination of high
46 scores on the insecurity and proximity seeking sub-scales of the VASQ) showed greater
47 stress-induced cortisol reactivity than participants with secure attachment style. Individuals
48 with avoidant attachment style (high scores for insecurity and low scores for proximity
49 seeking) did not differ from the secure attachment style group in terms of their cortisol 11
50
51
52
53
54
55
56
57
58
59
60

1
2
3 reactivity. Attachment style was not associated with the post-stress recovery in cortisol
4 concentrations. These data provide evidence for an association between attachment style
5 and increased reactivity of the HPA axis in response to a standardised group psychosocial
6 stressor in healthy young female participants.
7
8
9

10
11
12
13 The results are consistent with the work of Quirin et al. (2008), which also showed that
14 anxious attachment style predicted greater cortisol responding in females with no effect for
15 avoidant attachment. However, the results are in contrast with other work showing that
16 avoidant (but not anxious) attachment predicted enhanced stress-induced cortisol
17 responding in females (Powers et al., 2006). The results are also inconsistent with other
18 studies showing no relationship between attachment style and cortisol reactivity (Ditzen et
19 al., 2008; Smeets, 2010).
20
21
22
23
24
25
26
27
28

29 These discrepancies in findings may in part be attributable to the choice of stressor. We chose
30 to use an adapted form of the TSST for use with groups. The TSST is a reliable activator of
31 HPA axis function comprising the key elements of uncontrollability and socio-evaluative threat
32 (Dickerson et al., 2004). It is not possible to compare the size of the cortisol response described
33 here to those that would be elicited by the individual TSST; hence
34 whether the group nature of the stressor represented a particularity potent stimulus for this
35 group is undecided. It would be interesting to repeat this study using the individual TSST to
36 explore further this possibility. However, it is noteworthy that the modifications to the TSST-
37 G employed here (i.e. free group interactions in the preparatory phase and open visibility
38 during the stressor) produced a statistically significant cortisol stress reactivity response.
39 This provides opportunities for exploration of other social interventions in stress responding
40 such as reported by Häusser et al. (2012) in terms of group social identity.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Another distinguishing feature of this study was the use of the VASQ for the assessment of
4 attachment style. We chose to use this self-report tool as it has been shown to perform
5 somewhat better in predicting depression than other self-report measures of attachment
6 (Bifulco et al., 2003), as well as predicting negative psychosocial well-being and mental
7 health in university students (Carr et al., 2013). The study has several other strengths in that
8 self-reported menstrual phase, age, smoking and BMI were all accounted for in the
9 modelling of cortisol reactivity. The study also controlled for time of day and collected
10 multiple saliva samples at 10 minute intervals for more than 60 minutes, providing a full
11 neuroendocrine response profile, enabling accurate examination of individual cortisol
12 reactivity and recovery.
13
14
15
16
17
18
19
20
21
22
23
24

25 The findings may reflect subtle underlying changes in HPA axis function linked to attachment
26 style that are important for future mental health and well-being. For example an enhanced
27 cortisol response to the TSST has been shown to predict depressive symptoms in young adults
28 (Morris et al., 2012) and suicidal ideation in female adolescents with a history of mental health
29 concerns (Giletta et al., 2014). The results are also consistent with evidence showing greater
30 cortisol reactivity to the TSST in older adults subjected to separation from both parents during
31 childhood (Pesonen et al., 2010) and in young adults exposed to
32 severe pre-natal stress (Entringer et al., 2009).
33
34
35
36
37
38
39
40
41
42
43

44 The findings may also inform basal HPA axis function as it has been shown that the cortisol
45 response to laboratory stress is positively associated with average cortisol concentrations
46 over the day (Kidd et al., 2014). This may provide an eventual route to allostatic overload
47 and negative physical and mental health outcomes (McEwen, 2000; Morris et al., 2012). It
48 may also underpin observed aberrant diurnal profiles of cortisol secretion in anxious
49 attachment style (Oskis et al., 2011; Quirin et al., 2008). The study also provided supportive
50 evidence that avoidant insecure attachment style is somewhat similar to secure attachment
51
52
53
54
55
56
57
58
59
60

1
2
3 in terms of neuroendocrine function, consistent with evidence concerning the cortisol
4 awakening response (Oskis et al., 2011), as well as the findings that there are lower health
5 risks in insecure avoidant individuals compared to those anxiously attached (Bifulco et al.,
6 2002a; Fraley et al., 2004; Sbarra et al., 2013).
7
8
9
10

11
12
13 The current results are limited to healthy young females so it would be interesting to repeat
14 the study in healthy young males and with different age ranges. Another limitation is the
15 reliance upon self-reported menstrual phase, not hormonal assessment. Also the cross-
16 sectional design, with no long-term follow-up in relation to health outcomes, means we are
17 unable to draw any conclusions about whether the observed results are implicated in future
18 health outcomes.
19
20
21
22
23
24

25 26 27 **Conclusions**

28
29 In conclusion the study used an adapted version of the newly developed TSST-G to explore the
30 impact of attachment style on acute stress responding within a group setting. Healthy young
31 females with anxious attachment style showed a more marked cortisol response to the stressor.
32 Results obtained were not related to age, self-reported menstrual phase, smoking status or
33 BMI. The results indicate that differences in HPA axis activation may
34 provide a pre-clinical indication of ill-health vulnerability.
35
36
37
38
39
40
41
42

43 **Acknowledgements**

44 We would like to acknowledge the Bial Foundation, who funded this study.
45
46
47

48 **Declaration of Interest**

49 The authors report no conflicts of interest
50
51
52
53
54
55
56
57
58
59
60

References

- 1
2
3
4
5
6
7 Bifulco A, Mahon J, Kwon JH, Moran PM, Jacobs C. (2003). The vulnerable attachment
8 style questionnaire (VASQ): an interview-based measure of attachment styles that
9 predict depressive disorder. *Psychol Med* 33(06): 1099-110.
- 10
11 Bifulco A, Moran PM, Ball C, Bernazzani O. (2002a). Adult attachment style. I: Its
12 relationship to clinical depression. *Soc Psych Psych Epid* 37(2): 50-59.
- 13
14 Bifulco A, Moran PM, Ball C, Lillie A. (2002b). Adult attachment style. II: Its relationship to
15 psychosocial depressive-vulnerability. *Soc Psych Psych Epid* 37(2): 60-67.
- 16
17 Brennan KA, Clark CL, Shaver PR. (1998). Self-report measurement of adult attachment
18 *Attachment theory and close relationships* (pp. 46-76). New York: Guilford Press.
- 19
20 Brennan KA, Shaver PR. (1995). Dimensions of adult attachment, affect regulation, and
21 romantic relationship functioning *Pers Soc Psychol B* 21(3): 267-83.
- 22
23 Burnette JL, Davis DE, Green JD, Worthington Jr EL, Bradfield E. (2009). Insecure
24 attachment and depressive symptoms: The mediating role of rumination, empathy,
25 and forgiveness. *Pers Individ Dif* 46(3): 276-80.
- 26
27 Carr S, Colthurst K, Coyle M, Elliott D. (2013). Attachment dimensions as predictors of
28 mental health and psychosocial well-being in the transition to university. *Eur J*
29 *Psychol Educ* 28(2): 157-72.
- 30
31 Diamond LM. (2001). Contributions of psychophysiology to research on adult attachment:
32 Review and recommendations. *Pers Soc Psychol Rev* 5(4): 276-95.
- 33
34 Dickerson SS, Kemeny ME. (2004). Acute stressors and cortisol responses: a theoretical
35 integration and synthesis of laboratory research. *Psychol Bull* 130(3): 355-91.
- 36
37 Ditzen B, Schmidt S, Strauss B, Nater UM, Ehlert U, Heinrichs M. (2008). Adult attachment
38 and social support interact to reduce psychological but not cortisol responses to
39 stress. *J Psychosom Res* 64(5): 479-86.
- 40
41 Dockray S, Susman EJ, Dorn LD. (2009). Depression, Cortisol Reactivity, and Obesity in
42 Childhood and Adolescence. *J Adolesc Health* 45(4): 344-50.
- 43
44 Entringer S, Kumsta R, Hellhammer DH, Wadhwa PD, Wüst S. (2009). Prenatal exposure
45 to maternal psychosocial stress and HPA axis regulation in young adults. *Horm*
46 *Behav* 55(2): 292-98.
- 47
48 Fraley RC, Bonanno GA. (2004). Attachment and loss: A test of three competing models on
49 the association between attachment-related avoidance and adaptation to
50 bereavement. *Pers Soc Psychol B* 30(7): 878-90.
- 51
52
53
54
55
56
57
58
59
60

- 1
2
3 Giletta M, Calhoun C, Hastings P, Rudolph K, Nock M, Prinstein M. (2014). Multi-Level
4 Risk Factors for Suicidal Ideation Among at-Risk Adolescent Females: The Role of
5 Hypothalamic-Pituitary-Adrenal Axis Responses to Stress. *J Abnorm Child Psychol*:
6 1-14.
7
8
9 Häusser JA, Kattenstroth M, van Dick R, Mojzisch A. (2012). "We" are not stressed: Social
10 identity in groups buffers neuroendocrine stress reactions. *J Exp Soc Psychol* 48(4):
11 973-77.
12
13 Jinyao Y, Xiongzha Z, Auerbach RP, Gardiner CK, Lin C, Yuping W, Shuqiao Y. (2012).
14 Insecure attachment as a predictor of depressive and anxious symptomology.
15 *Depress Anxiety* 29(9): 789-96.
16
17
18 Kidd T, Carvalho LA, Steptoe A. (2014). The relationship between cortisol responses to
19 laboratory stress and cortisol profiles in daily life. *Biol Psychol* 99(0): 34-40.
20
21 Kidd T, Hamer M, Steptoe A. (2011). Examining the association between adult attachment
22 style and cortisol responses to acute stress. *Psychoneuroendocrinology* 36(6): 771-
23 79.
24
25
26 Kiecolt-Glaser JK, Newton T, Cacioppo JT, MacCallum RC, Glaser R, Malarkey WB.
27 (1996). Marital conflict and endocrine function: Are men really more physiologically
28 affected than women? *J Consult Clin Psychol* 64(2): 324.
29
30
31 Kirschbaum C, Klauer T, Filipp S-H, Hellhammer DH. (1995). Sex-specific effects of social
32 support on cortisol and subjective responses to acute psychological stress.
33 *Psychosom Med* 57(1): 23-31.
34
35
36 Kirschbaum C, Pirke KM, Hellhammer DH. (1993). The 'Trier Social Stress Test'--a tool for
37 investigating psychobiological stress responses in a laboratory setting.
38 *Neuropsychobiology* 28(1-2): 76-81.
39
40
41 McEwen BS. (2000). Allostasis and allostatic load: implications for
42 neuropsychopharmacology. *Neuropsychopharmacology* 22(2): 108-24.
43
44
45 Morris MC, Rao U, Garber J. (2012). Cortisol responses to psychosocial stress predict
46 depression trajectories: Social-evaluative threat and prior depressive episodes as
47 moderators. *J Affect Disord* 143(1): 223-30.
48
49
50 Oskis A, Clow A, Loveday C, Hucklebridge F, Sbarra D. (2014). Biological Stress Regulation in
51 Female Adolescents: A Key Role for Confiding. *J Youth Adolescence*: 1-12.
52
53
54 Oskis A, Loveday C, Hucklebridge F, Thorn L, Clow A. (2011). Anxious attachment style
55 and salivary cortisol dysregulation in healthy female children and adolescents. *J*
56 *Child Psychol Psyc* 52(2): 111-18.
57
58
59
60 Pesonen A-K, Räikkönen K, Feldt K, Heinonen K, Osmond C, Phillips DIW, Barker DJP, et
al. (2010). Childhood separation experience predicts HPA axis hormonal responses 16

- 1
2
3 in late adulthood: A natural experiment of World War II. *Psychoneuroendocrinology*
4 35(5): 758-67.
5
6 Pierrehumbert B, Torrisi R, Ansermet F, Borghini A, Halfon O. (2012). Adult attachment
7 representations predict cortisol and oxytocin responses to stress. *Attach Hum Dev*
8 14(5): 453-76.
9
10 Powers SI, Pietromonaco PR, Gunlicks M, Sayer A. (2006). Dating couples' attachment
11 styles and patterns of cortisol reactivity and recovery in response to a relationship
12 conflict. *J Pers Soc Psychol* 90(4): 613.
13
14 Puig J, Englund MM, Simpson JA, Collins WA. (2013). Predicting adult physical illness from
15 infant attachment: A prospective longitudinal study. *Health Psychol* 32(4): 409.
16
17 Quirin M, Pruessner JC, Kuhl J. (2008). HPA system regulation and adult attachment
18 anxiety: Individual differences in reactive and awakening cortisol.
19
20 *Psychoneuroendocrinology* 33(5): 581-90.
21
22 Repetti RL, Taylor SE, Seeman TE. (2002). Risky families: family social environments and
23 the mental and physical health of offspring. *Psychol Bull* 128(2): 330.
24
25 Rifkin-Graboi A. (2008). Attachment status and salivary cortisol in a normal day and during
26 simulated interpersonal stress in young men. *Stress* 11(3): 210-24.
27
28 Sbarra DA, Borelli JL. (2013). Heart rate variability moderates the association between
29 attachment avoidance and self-concept reorganization following marital separation.
30
31 *Int J Psychophysiol* 88(3): 253-60.
32
33 Shaver PR, Mikulincer M. (2002). Attachment-related psychodynamics. *Attach Hum Dev*
34 4(2): 133-61.
35
36 Smeets T. (2010). Autonomic and hypothalamic–pituitary–adrenal stress resilience: Impact
37 of cardiac vagal tone. *Biol Psychol* 84(2): 290-95.
38
39 Smyth N, Hucklebridge F, Thorn L, Evans P, Clow A. (2013). Salivary cortisol as a
40 biomarker in social science research. *Soc Personal Psychol Compass* 7(9): 605-25.
41
42 Stroud LR, Salovey P, Epel ES. (2002). Sex differences in stress responses: social
43 rejection versus achievement stress. *Biol Psychiat* 52(4): 318-27.
44
45 von Dawans B, Kirschbaum C, Heinrichs M. (2011). The Trier Social Stress Test for
46 Groups (TSST-G): A new research tool for controlled simultaneous social stress
47 exposure in a group format. *Psychoneuroendocrinology* 36(4): 514-22.
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1 Mean (\pm S.E.M.) salivary free cortisol concentrations (nmol/l) for all participants (N = 78). A: immediately before onset of the TSST-G; B mid-way through the TSST-G; C immediately after the end of the TSST-G.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2 Attachment style differences in cortisol reactivity. Reactivity was significantly greater for the insecure anxious attachment style group in comparison to the securely ($p=0.011$).

Table 1 Descriptive statistics and intercorrelations between cortisol, VASQ attachment measures and demographic data (N=78)

Variables	Descriptives	Correlations							
		Cortisol recovery	Vulnerability	Insecurity	Proximity	Age	Menstrual cycle phase	Smoking status	BMI
Cortisol reactivity M (SD)	4.47 (6.50)	.185	.289*	.269*	.177	.349**	-.141	-.103	.011
Cortisol recovery M (SD)	4.20 (4.50)		-.020	-.047	.016	.276*	-.176	.015	.017
VASQ Vulnerability M (SD)	60.98 (9.81)			.775**	.770**	.209	-.037	.163	-.119
VASQ Insecurity M (SD)	33.37 (6.39)				.193	.247*	-.051	.001	-.035
VASQ Proximity M (SD)	27.61 (6.32)					-.075	-.006	.254*	-.150
Age M (SD)	20.22 (3.21)						-.054	.099	.218
Menstrual cycle phase % luteal	38.5							-.017	-.007
Smoking status % non-smoker	85.9								-.111
BMI M (SD)	21.46 (3.76)								

* $p < .05$, ** $p < .001$

Table 2 Prediction of cortisol reactivity

Predictors	beta	<i>t</i>	<i>p</i>
VASQ Vulnerability	.241	2.205	.031
Age	.322	2.902	.005
Menstrual cycle phase	-.118	-1.132	.261
Smoking status	-.182	-1.706	.092
BMI	-.051	-.468	.641
R^2	.162 ($p = 0.003$)		